

Clear



Registered Name: Qolt Siham Sahara

Nickname: Qolt

Multi-Drug Resistance 1, (MDR1)

Registration ID: HP55542103

Microchip: REDACTED
Breed: Sloughi
Gender: Male

Owner: Erika Wyatt

Country: United States

Testing date: 2020/7/18

### Test results - Known disorders in the breed

Disorder	Туре	Mode of Inheritance	Result
Rod-Cone Dysplasia 1a, (rdc1a); mutation originally found in Sloughi	Ocular Disorders	Autosomal Recessive	Clear
Test results for pharmacogenetics			
Disorder		Mode of Inheritance	Result

On behalf of Genoscoper Laboratories,

SIGNATURE

Jonas Donner, PhD, Head of Research and Development at Genoscoper Laboratories

**Autosomal Dominant** 





Nickname: Qolt

Registration ID: HP55542103

Microchip: 941000019903816

Breed: Sloughi
Gender: Male

Owner: Erika Wyatt

Country: United States

Testing date: 2020/7/18

### Test results - Traits - page 1

### **Coat Type**

Trait	Genotype	Description
Coat Length	L/L	The dog is likely to have short-haired coat.
Furnishings / Improper Coat in Portuguese Water Dogs (marker test)	GG/CC	The dog is not genetically likely to express furnishings.
KRT71 c.451C>T (p.Arg151Trp)	C/C	The dog does not carry any copies of the tested allele causing curly coat. The dog most likely has non-curly hair.
MC5R c.237A>T	C/T	The dog carries one copy of the allele associated with heavy shedding and one copy of the allele associated low shedding. This genotype has no effect on a dog with furnishings, but non-wire-haired dog with this genotype is likely heavy or seasonal shedder.
SGK3 (p.Val96Glyfs)	1/1	The dog does not carry the tested hairlessness allele of the American Hairless Terrier.
SGK3 c.137_138insT (p.Glu47Glyfs)	D/D	The dog does not carry the tested hairlessness allele of the Scottish Deerhound.

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Qolt

Registration ID: HP55542103

Microchip: 941000019903816

Breed: Sloughi
Gender: Male

Owner: Erika Wyatt

Country: United States

**Testing date:** 2020/7/18

### Test results - Traits - page 2

#### **Coat Colour**

Trait	Genotype	Description
Colour Locus E - Extensions	Em/E	The dog is likely to have a dark mask.
Colour Locus B - Brown	B/B	The dog is not likely to have brown pigment.
Colour Locus K - Dominant Black	KB/ky    KB/kbr    kbr/ky    kbr/kbr	The dog is genetically dominant black or brindle.
Colour Locus A - Agouti	ay/at	The dog is genetically sable. The dog carries tan points or saddle tan colour.
Colour Locus S - Piebald or extreme white spotting	S/S	The dog is likely to have solid coat colour with minimal white.
Colour Locus H - Harlequin	h/h	The dog doesn't have harlequin pattern.
Dilution (d <sup>2</sup> allele)	D/D	The dog does not carry any copies of the rare d2 allele associated with dilution in Chow Chow, French Bulldog, Sloughi and Thai Ridgeback.
Merle (M allele)	m/m	The dog is genetically non-merle and does not carry a <i>SILV</i> gene SINE insertion.
Saddle Tan (RALY gene dupl.)	-/dup	The dog may have saddle tan pattern if it has also tan point genotype at the A locus.
Albinism (caL-allele)	C/C	The dog does not carry the tested mutation for albinism.

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Qolt

Registration ID: HP55542103

Microchip: 941000019903816

Breed: Sloughi
Gender: Male

Owner: Erika Wyatt

Country: United States

**Testing date:** 2020/7/18

### Test results - Traits - page 3

#### **Body Size**

Trait	Genotype	Description
IGF1 (chr15:41221438)	G/G	The dog is homozygous for the ancestral allele typically associated with large body mass.
IGF1R c.611G>A (p.Arg204His)	G/G	The dog carries two ancestral alleles typically found in larger-sized breeds.
ACSL4 chrX.82919525C>T	C/C	The dog doesn't have the allele associated with large skeletal size and heavy muscling with considerable back fat thickness.
IGSF1 p.Asp768Glu	A/A	The dog has two copies of the allele associated with heavy muscling.
IRS4 chrX:82296039	G/G	The dog doesn't have the allele associated with large body size.
FGF4 insertion	D/D	The dog is homozygous for the ancient allele. The dog is likely to have legs of normal length.
STC2 (chr4:39182836)	T/T	The dog has two copies of the ancestral allele associated with larger body size.
GHR1 (p.Glu191Lys)	G/G	The dog has two copies of the ancestral allele associated with larger body size.
GHR2 (p.Pro177Leu)	C/C	The dog has two copies of the ancestral allele associated with larger body size.
HMGA2 (chr10:8348804)	G/G	The dog has two copies of the ancestral allele associated with larger body size.

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Qolt

Registration ID: HP55542103

Microchip: 941000019903816

Breed: Sloughi
Gender: Male

Owner: Erika Wyatt
Country: United States

**Testing date:** 2020/7/18

### Test results - Traits - page 4

### Morphology

Trait	Genotype	Description
BMP3 c.1344C>A (p.Phe448Leu)	C/C	The dog does not carry the tested allele typically associated with shortened head (brachycephaly). The dog is more likely to have an elongated head (dolichocephaly).
SMOC2	D/D	The dog does not carry the tested allele typically associated with shortened head (brachycephaly). The dog is more likely to have an elongated head (dolichocephaly).
chr10:11072007	C/C	The dog carries two copies of an allele typically associated with floppy ears. The dog is more likely to have floppy than pricked ears.
T c.189C>G (p.lle63Met)	C/C	The dog does not carry the tested bobtail-causing genetic variant. The dog is most likely long-tailed.
EPAS1 (p.Gly305Ser)	G/G	The dog does not carry the tested variant associated with adaptation to high altitudes.
LIMBR1 DC-1	G/G	The dog does not carry the tested allele associated with hind dewclaws in Asian breeds. The dog is not likely to have hind dewclaws.
LIMBR1 DC-2	G/G	The dog does not carry the tested allele associated with hind dewclaws in western breeds. The dog is likely not to have hind dewclaws.
AXL4	D/D	The dog does not have the tested allele typically associated with blue eyes in Siberian Huskies. The dog is likely to have brown eyes.

On behalf of Genoscoper Laboratories,

SIGNATURE



### **Blood Disorders - page 1**

Bleeding disorder due to P2RY12 defect Canine Cyclic Neutropenia, Cyclic Hematopoiesis, Grey Collie Syndrome, Autosomal Recessive Clear Canine Leukocyte Adhesion Deficiency (CLAD), type III Autosomal Recessive Clear Canine Scott Syndrome, (CSS) Autosomal Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier Factor IX Deficiency or Hemophilia B; mutation originally found in Lasa Ayso Factor VII Deficiency or Hemophilia B; mutation originally found in Boxer Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd Dog Factor VIII Deficiency or Hemophilia A; mutation originally found in Havanese Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Havanese Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Havanese Clear Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VII Deficiency Autosomal Dominant Factor VII Deficiency Clear Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pomeranian Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pomeranian Autosomal Recessive Clear Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pomeranian Autosomal Recessive Clear Hereditary Eliptocytosis Clear Hereditary Plosphofructokinase (PFK) Deficiency Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear	Disorder	Mode of Inheritance	Result
CAIN Canine Leukocyte Adhesion Deficiency (CLAD), type III Autosomal Recessive Clear Canine Scott Syndrome, (CSS) Autosomal Recessive Clear Factor IX Deficiency or Hemophilia B; mutation Gly379Glu X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Soure X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.C.y.s548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency or Hemophilia A; p.C.y.s548Tyr mutation originally found in Old English Sheepdog Factor XI Deficiency or Hemophilia A; p.C.y.s548Tyr mutation originally Autosomal Dominant (Incomplete Penetrance) Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance) Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Hereditary Elliptocytosis Clear Hereditary Elliptocytosis Clear Hereditary Phosphofructokinase (PFK) Deficiency Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear	Bleeding disorder due to P2RY12 defect	Autosomal Recessive	Clear
Canine Scott Syndrome, (CSS) Autosomal Recessive Clear Factor IX Deficiency or Hemophilia B; mutation Gly379Glu X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance) Familial Congenital Methemoglobinemia; mutation originally found in Autosomal Recessive Clear Factor XI Deficiency Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Factor XI Deficiency Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Factor XI Deficiency Clear Factor XI Deficiency Autosomal Recessive Clear Factor XI Deficiency Clear Factor XI Deficiency Autosomal Recessive Clear		Autosomal Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation Gly379Glu X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor VII Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in Factor XI Deficiency  Factor XI Deficiency  Autosomal Recessive  Clear  Clear  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk  Autosomal Recessive  Clear	Canine Leukocyte Adhesion Deficiency (CLAD), type III	Autosomal Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier  Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Autosomal Recessive Clear Apso  Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Cerman Shepherd Dog Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Havanese Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Havanese Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old X-linked Recessive Clear English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Clear Indicate Penetrance Clear Indicate Penetrance Pomeranian Clear Indicate Penetrance Penetran	Canine Scott Syndrome, (CSS)	Autosomal Recessive	Clear
Terrier Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Autosomal Recessive Clear Factor VIII Deficiency Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance) Familial Congenital Methemoglobinemia; mutation originally found in Autosomal Recessive Clear Premean Mountain Dog Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Pyrenean Mountain Dog Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Morfolk Premeditary Elliptocytosis Clear Hereditary Phosphofructokinase (PFK) Deficiency Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear	Factor IX Deficiency or Hemophilia B; mutation Gly379Glu	X-linked Recessive	Clear
Apso Factor VII Deficiency Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Factor VIII Deficiency or Hemophilia A; mutation originally found in Havanese Factor VIII Deficiency or Hemophilia A; mutation originally found in Havanese Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance) Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mautosomal Recessive Clear Clear Hereditary Elliptocytosis Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear		X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance) Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Pyrenean Mountain Dog Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive Clear Hereditary Elliptocytosis Clear Hereditary Phosphofructokinase (PFK) Deficiency Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim Terrier		X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd Dog  Factor VIII Deficiency or Hemophilia A; mutation originally found in Havanese  Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally English Sheepdog  Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance)  Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Hereditary Elliptocytosis  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim Terrier  Clear	Factor VII Deficiency	Autosomal Recessive	Clear
German Shepherd Dog  Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear  Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear  Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance)  Familial Congenital Methemoglobinemia; mutation originally found in Autosomal Recessive Clear  Pomeranian  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear  Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs  Hereditary Elliptocytosis Clear  Hereditary Phosphofructokinase (PFK) Deficiency Autosomal Recessive Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier	Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer	X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive  Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive  Factor XI Deficiency  Autosomal Dominant (Incomplete Penetrance)  Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Hereditary Elliptocytosis  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier		X-linked Recessive	Clear
English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd  Factor XI Deficiency  Autosomal Dominant (Incomplete Penetrance)  Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Hereditary Elliptocytosis  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier		X-linked Recessive	Clear
Factor XI Deficiency  Autosomal Dominant (Incomplete Penetrance)  Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Hereditary Elliptocytosis  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier		X-linked Recessive	Clear
Familial Congenital Methemoglobinemia; mutation originally found in Pomeranian  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Hereditary Elliptocytosis  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier		X-linked Recessive	Clear
Pomeranian  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive  Clear  Hereditary Elliptocytosis  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  Clear	Factor XI Deficiency		Clear
Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs  Hereditary Elliptocytosis  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  Clear		Autosomal Recessive	Clear
mixed breed dogs  Hereditary Elliptocytosis  Clear  Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  Clear		Autosomal Recessive	Clear
Hereditary Phosphofructokinase (PFK) Deficiency  Autosomal Recessive  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  Clear		Autosomal Recessive	Clear
Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear and Cairn Terrier	Hereditary Elliptocytosis		Clear
and Cairn Terrier	Hereditary Phosphofructokinase (PFK) Deficiency	Autosomal Recessive	Clear
May-Hegglin Anomaly (MHA) Autosomal Dominant Clear		Autosomal Recessive	Clear
	May-Hegglin Anomaly (MHA)	Autosomal Dominant	Clear



### **Blood Disorders - page 2**

Disorder	Mode of Inheritance	Result
Prekallikrein Deficiency	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Basenji	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Beagle	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Pug	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear
Trapped Neutrophil Syndrome, (TNS)	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 1	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 2	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Kooikerhondje	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Scottish Terrier	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear



### Ocular Disorders - page 1

Disorder	Mode of Inheritance	Result
Canine Multifocal Retinopathy 1, (CMR1); mutation originally found in Mastiff-related breeds	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 2, (CMR2); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 3, (CMR3); mutation originally found in Lapponian Herder	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shepherd Dog	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shorthaired Pointer	Autosomal Recessive	Clear
Cone-Rod Dystrophy 1, (crd1); mutation originally found in American Staffordshire Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy 2, (crd2); mutation originally found in American Pit Bull Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy, (cord1-PRA / crd4)	Autosomal Recessive (Incomplete Penetrance)	Clear
Cone-Rod Dystrophy, Standard Wirehaired Dachshund, (crd SWD)	Autosomal Recessive	Clear
Congenital Eye Disease; mutation originally found in Irish Soft-Coated Wheaten Terrier	Autosomal Recessive	Clear
Dominant Progressive Retinal Atrophy, (DPRA)	Autosomal Dominant	Clear
Early Onset PRA (EOPRA); mutation originally found in Portuguese Water Dog	Autosomal Recessive	Clear
Early Retinal Degeneration, (erd); mutation originally found in Norwegian Elkhound	Autosomal Recessive	Clear
Generalized Progressive Retinal Atrophy	Autosomal Recessive	Clear
Golden Retriever Progressive Retinal Atrophy 1, (GR_PRA 1)	Autosomal Recessive	Clear
Goniodysgenesis and glaucoma; mutation originally found in Border Collie	Autosomal Recessive	Clear
Italian Greyhound Progressive Retinal Atrophy 1 (IG-PRA1)	Autosomal Recessive	Clear
Primary Hereditary Cataract, (PHC); mutation originally found in Australian Shepherd	Autosomal Dominant (Incomplete Penetrance)	Clear
Primary Lens Luxation, (PLL)	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Basset Fauve de Bretagne	Autosomal Recessive	Clear



### Ocular Disorders - page 2

Disorder	Mode of Inheritance	Result
Primary Open Angle Glaucoma, (POAG); mutation originally found in Beagle	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Norwegian Elkhound	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Petit Basset Griffon Vendeen	Autosomal Recessive	Clear
Primary lens luxation (PLL) and glaucoma; mutation originally found in Shar Pei	Autosomal Recessive	Clear
Progressive Retinal Atrophy (PRA4); mutation originally found in Lhasa Apso	Autosomal Recessive	Clear
Progressive Retinal Atrophy Type III, (PRA type III); mutation originally found in Tibetan Spaniel and Tibetan Terrier	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (CNGA1-PRA); mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PAP1_PRA); mutation originally found in Papillon and Phalene	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Basenji	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Swedish Vallhund	Autosomal Recessive	Clear
Rod-Cone Dysplasia 1, (rcd1); mutation originally found in Irish Setter	Autosomal Recessive	Clear
Rod-Cone Dysplasia 3, (rcd3)	Autosomal Recessive	Clear
X-Linked Progressive Retinal Atrophy 1, (XLPRA1)	X-linked Recessive	Clear
X-Linked Progressive Retinal Atrophy 2, (XLPRA2; Type A PRA)	X-linked Recessive	Clear

### **Cardiac Disorders**

Disorder	Mode of Inheritance	Result
Dilated Cardiomyopathy, (DCM); mutation originally found in Schnauzer	Autosomal Recessive	Clear
Long QT Syndrome	Autosomal Dominant	Clear



#### **Endocrine Disorders**

Disorder	Mode of Inheritance	Result
Congenital Dyshormonogenic Hypothyroidism with Goiter; mutation originally found in Shih Tzu	Autosomal Recessive	Clear
Congenital Hypothyroidism; mutation originally found in Tenterfield Terrier	Autosomal Recessive	Clear
Congenital Hypothyroidism; mutation originally found in Toy Fox and Rat Terrier	Autosomal Recessive	Clear

#### **Immunological Disorders**

Disorder	Mode of Inheritance	Result
Autosomal Recessive Severe Combined Immunodeficiency, (ARSCID)	Autosomal Recessive	Clear
Complement 3 (C3) Deficiency	Autosomal Recessive	Clear
Myeloperoxidase Deficiency	Autosomal Recessive	Clear
Severe Combined Immunodeficiency in Frisian Water Dogs, (SCID)	Autosomal Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Basset Hound	X-linked Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Cardigan Welsh Corgi	X-linked Recessive	Clear



### **Renal Disorders**

Disorder	Mode of Inheritance	Result
2,8-Dihydroxyadenine (2,8-DHA) urolithiasis	Autosomal Recessive	Clear
Cystic Renal Dysplasia and Hepatic Fibrosis; mutation originally found in Norwich Terrier	Autosomal Recessive	Clear
Cystinuria Type I-A; mutation originally found in Newfoundland Dog	Autosomal Recessive	Clear
Cystinuria Type II-A; mutation originally found in Australian Cattle Dog	Autosomal Dominant	Clear
Fanconi Syndrome	Autosomal Recessive	Clear
Hyperuricosuria, (HUU)	Autosomal Recessive	Clear
Polycystic Kidney Disease in Bull Terriers, (BTPKD)	Autosomal Dominant	Clear
Primary Hyperoxaluria, (PH); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Protein Losing Nephropathy, (PLN); NPHS1 gene variant		Clear
Renal Cystadenocarcinoma and Nodular Dermatofibrosis, (RCND)	Autosomal Dominant	Clear
X-Linked Hereditary Nephropathy, (XLHN)	X-linked Recessive	Clear
X-Linked Hereditary Nephropathy, (XLHN); mutation originally found in Navasota Dog	X-linked Recessive	Clear
Xanthinuria, Type 1a; mutation originally found in mixed breed dogs	Autosomal Recessive	Clear
Xanthinuria, Type 2a; mutation originally found in Toy Manchester Terrier	Autosomal Recessive	Clear
Xanthinuria, Type 2b; mutation originally found in Cavalier King Charles Spaniel and English Cocker Spaniel	Autosomal Recessive	Clear



#### **Metabolic Disorders**

Disorder	Mode of Inheritance	Result
Glycogen Storage Disease Type II or Pompe's Disease, (GSD II)	Autosomal Recessive	Clear
Glycogen Storage Disease Type IIIa, (GSD IIIa)	Autosomal Recessive	Clear
Glycogen Storage Disease Type Ia, (GSD Ia)	Autosomal Recessive	Clear
Hypocatalasia or Acatalasemia	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Beagle	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Border Collie	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in Dachshund	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in New Zealand Huntaway	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in Brazilian Terrier	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in German Shepherd	Autosomal Recessive	Clear
Pyruvate Dehydrogenase Phosphatase 1 (PDP1) Deficiency	Autosomal Recessive	Clear



#### **Muscular Disorders**

Disorder	Mode of Inheritance	Result
Cavalier King Charles Spaniel Muscular Dystrophy, (CKCS-MD)	X-linked Recessive	Clear
Centronuclear Myopathy, (CNM); mutation originally found in Great Dane	Autosomal Recessive	Clear
Centronuclear Myopathy, (CNM); mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Duchenne or Dystrophin Muscular Dystrophy, (DMD); mutation originally found in Golden Retriever	X-linked Recessive	Clear
Duchenne or Dystrophin Muscular Dystrophy, (DMD); mutation originally found in Norfolk Terrier	X-linked Recessive	Clear
Muscular Dystrophy, Ullrich-type; mutation originally found in Landseer	Autosomal Recessive	Clear
Myostatin deficiency (Double Muscling, "Bully")	Autosomal Recessive	Clear
Myotonia Congenita; mutation originally found in Australian Cattle Dog	Autosomal Recessive	Clear
Myotubular Myopathy; mutation originally found in Rottweiler	X-linked Recessive	Clear
Nemaline Myopathy; mutation originally found in American Bulldog	Autosomal Recessive	Clear
X-Linked Myotubular Myopathy	X-linked Recessive	Clear



### Neurological Disorders - page 1

Disorder	Mode of Inheritance	Result
Acral Mutilation Syndrome, (AMS)	Autosomal Recessive	Clear
Alaskan Husky Encephalopathy, (AHE)	Autosomal Recessive	Clear
Alexander Disease (AxD); mutation originally found in Labrador Retriever	Autosomal Dominant	Clear
Bandera's Neonatal Ataxia, (BNAt)	Autosomal Recessive	Clear
Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy	Autosomal Recessive	Clear
Cerebellar Cortical Degeneration, (CCD); mutation originally found in Vizsla	Autosomal Recessive	Clear
Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun	Autosomal Recessive	Clear
Dandy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier	Autosomal Recessive	Clear
Degenerative Myelopathy, (DM; SOD1A)	Autosomal Recessive (Incomplete Penetrance)	Clear
Early-Onset Progressive Polyneuropathy; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear
Fetal Onset Neuroaxonal Dystrophy, (FNAD)	Autosomal Recessive	Clear
Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter	Autosomal Recessive	Clear
Hereditary Ataxia; mutation originally found in in Norwegian Buhund	Autosomal Recessive	Clear
Hyperekplexia or Startle Disease	Autosomal Recessive	Clear
Hypomyelination; mutation originally found in Weimaraner	Autosomal Recessive	Clear
Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback	Autosomal Recessive	Clear
Juvenile encephalopathy; mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear
L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier	Autosomal Recessive	Clear
L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear
Lagotto Storage Disease, (LSD)	Autosomal Recessive	Clear
Neonatal Cerebellar Cortical Degeneration or Cerebellar Abiotrophy, (NCCD)	Autosomal Recessive	Clear



### Neurological Disorders - page 2

, ()	Autosomal Recessive Autosomal Recessive	Clear
Neuroaxonal Dystrophy (NAD); mutation originally found in Rottweiler	Autosomal Recessive	
		Clear
Neuroaxonal Dystrophy (NAD); mutation originally found in Spanish Water A Dog	Autosomal Recessive	Clear
Neuroaxonal Dystrophy, (NAD); mutation originally found in Papillon	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 1, (NCL1); mutation originally found in Dachshund	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 10, (NCL10); mutation originally found in American Bulldog	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Alpine Dachsbracke	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Australian Shepherd	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in English Setter	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis, (NCL7); mutation originally found in Chinese Crested Dog and Chihuahua	Autosomal Recessive	Clear
Polyneuropathy with ocular abnormalities and neuronal vacuolation, (POANV); mutation originally found in Black Russian Terrier	Autosomal Recessive	Clear
Progressive Early-Onset Cerebellar Ataxia; mutation originally found in Finnish Hound	Autosomal Recessive	Clear
Sensory Neuropathy; mutation originally found in Border Collie	Autosomal Recessive	Clear
Shaking Puppy Spongiform LeucoEncephaloMyelopathy, (SLEM); mutation originally found in Border Terrier	Autosomal Recessive	Clear
Spinocerebellar Ataxia with Myokymia and/or Seizures (SCA)	Autosomal Recessive	Clear
Spinocerebellar Ataxia/ Late-Onset Ataxia (SCA, LOA)	Autosomal Recessive	Clear
Spongy Degeneration with Cerebellar Ataxia, (SDCA1); mutation originally A found in Belgian Shepherd Dog	Autosomal Recessive	Clear
Spongy Degeneration with Cerebellar Ataxia, (SDCA2); mutation originally A found in Belgian Shepherd Dog	Autosomal Recessive	Clear
X-Linked Tremors; mutation originally found in English Springer Spaniel	X-linked Recessive	Clear



#### **Neuromuscular Disorders**

Disorder	Mode of Inheritance	Result
Congenital Myasthenic Syndrome (CMS); mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Jack Russell Terrier	Autosomal Recessive	Clear
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Old Danish Pointing Dog	Autosomal Recessive	Clear
Episodic Falling Syndrome, (EFS)	Autosomal Recessive	Clear
Exercise-Induced Collapse, (EIC)	Autosomal Recessive (Incomplete Penetrance)	Clear
GM1 Gangliosidosis; mutation originally found in Portuguese Water Dog	Autosomal Recessive	Clear
GM2 Gangliosidosis, mutation originally found in Japanese Chin	Autosomal Recessive	Clear
GM2 Gangliosidosis; mutation originally found in Toy Poodle	Autosomal Recessive	Clear
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Irish Setter	Autosomal Recessive	Clear
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Terriers	Autosomal Recessive	Clear
Paroxysmal Dyskinesia, (PxD); mutation originally found in Irish Soft Coated Wheaten Terrier	Autosomal Recessive	Clear



#### **Skeletal Disorders**

Disorder	Mode of Inheritance	Result
Chondrodysplasia; mutation originally found in Norwegian Elkhound and Karelian Bear Dog	Autosomal Recessive	Clear
Cleft Palate; Cleft Lip and Palate with Syndactyly; ADAMTS20 gene mutation originally found in Nova Scotia Duck Tolling Retriever	Autosomal Recessive	Clear
Cleft Palate; DLX6 gene mutation originally found in Nova Scotia Duck Tolling Retriever	Autosomal Recessive	Clear
Craniomandibular Osteopathy, (CMO); mutation associated with terrier breeds	Autosomal Dominant (Incomplete Penetrance)	Clear
Hereditary Vitamin D-Resistant Rickets, (HVDRR)	Autosomal Recessive	Clear
Osteochondrodysplasia; mutation originally found in Miniature Poodle	Autosomal Recessive	Clear
Osteochondromatosis; mutation originally found in American Staffordshire Terrier	Autosomal Dominant	Clear
Osteogenesis Imperfecta, (OI); mutation originally found in Beagle	Autosomal Dominant	Clear
Osteogenesis Imperfecta, (OI); mutation originally found in Dachshund	Autosomal Recessive	Clear
Skeletal Disease (Hypophosphatasia); mutation originally found in Karelian Bear Dog	Autosomal Recessive	Clear
Skeletal Dysplasia 2, (SD2)	Autosomal Recessive	Clear
Spondylocostal Dysostosis	Autosomal Recessive	Clear
Van den Ende-Gupta Syndrome, (VDEGS)	Autosomal Recessive	Clear



#### **Dermal Disorders**

Disorder	Mode of Inheritance	Result
Dystrophic Epidermolysis Bullosa; mutation originally found in Central Asian Ovcharka	Autosomal Recessive	Clear
Dystrophic Epidermolysis Bullosa; mutation originally found in Golden Retriever	Autosomal Recessive	Clear
Epidermolytic Hyperkeratosis	Autosomal Recessive	Clear
Focal Non-Epidermolytic Palmoplantar Keratoderma, (FNEPPK); mutation originally found in Dogue de Bordeaux	Autosomal Recessive	Clear
Hereditary Footpad Hyperkeratosis, (HFH)	Autosomal Recessive	Clear
Hereditary Nasal Parakeratosis, (HNPK); mutation originally found in Greyhound	Autosomal Recessive	Clear
Ichthyosis; mutation originally found in American Bulldog	Autosomal Recessive	Clear
Ichthyosis; mutation originally found in Great Dane	Autosomal Recessive	Clear
Lamellar Ichthyosis, (LI)	Autosomal Recessive	Clear
Lethal Acrodermatitis, (LAD); mutation originally found in in Bull Terrier and Miniature Bull Terrier	Autosomal Recessive	Clear
Ligneous Membranitis	Autosomal Recessive	Clear
Musladin-Lueke syndrome, (MLS)	Autosomal Recessive	Clear
X-Linked Ectodermal Dysplasia, (XHED)	X-linked Recessive	Clear
<u> </u>		



#### **Other Disorders**

Disorder	Mode of Inheritance	Result
Acute Respiratory Distress Syndrome, (ARDS); mutation originally found in Dalmatian	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Italian Greyhound	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear
Congenital Keratoconjunctivitis Sicca and Ichthyosiform Dermatosis, (CKCSID)	Autosomal Recessive	Clear
Dental Hypomineralisation; mutation originally found in Border Collie	Autosomal Recessive	Clear
Lung Developmental Disease; mutation originally found in in Airedale Terrier	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Dachshund	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Persistent Müllerian Duct Syndrome, (PMDS); mutation originally found in Miniature Schnauzer	Autosomal Recessive	Clear
Primary Ciliary Dyskinesia, (PCD)	Autosomal Recessive	Clear





#### **APPENDIX**

### Explanation of the results of the tested disorders

Autosomal recessive inheritance (ARI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - A dog carries one copy of the tested mutation. Carriers typically have a normal, healthy appearance but pass on the mutation to approximately 50% of their offspring.

At risk - A dog carries two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

Autosomal dominant inheritance (ADI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

At risk - A dog carries one or two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

X-linked recessive inheritance (X-linked)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - Female carriers typically have a normal, healthy appearance but carry one copy of the tested mutation on one of their X chromosomes. As males only have one X chromosome, there are no male carriers.

At risk - Female dogs at risk carry two mutated copies of the tested mutation. Males carry one copy of the tested mutation on their single X chromosome. Dogs at risk are at high or increased risk of developing the disease/condition.

Please note that the descriptions above are generalized based on typically observed inheritance patterns. When obtaining a 'carrier' or 'at risk' test result, always refer to the corresponding online test documentation for more detailed information on the condition and any exceptions.





### Terms and Conditions

block-terms-and-conditions